



辣根过氧化物酶标记的山羊抗小鼠 IgG (γ 链特异性) Goat Anti-Mouse IgG (γ chain specific)/HRP

产品编号	规格	产品简介
ANM02-3	100ul	辣根过氧化物酶标记抗体广泛应用于免疫学、免疫细胞化学、分子生物学、病理学等领域。该产品为辣根过氧化物酶标记的经亲和层析的山羊抗小鼠IgG抗血清。经ELISA实验验证其与小鼠IgG ₁ , IgG _{2a} , IgG _{2b} , IgG ₃ 反应，与人免疫球蛋白有微量交叉反应。

应用范围

- 酶联免疫吸附实验(ELISA)
- 荧光免疫吸附实验(FLISA)
- 免疫印迹(immunoblotting)

稀释比例

推荐稀释比例 ELISA: 1:4000-1:8000, 具体稀释比例请根据实验确定。其他应用请根据具体实验确定最佳稀释度。

保存

HRP标记的山羊抗小鼠IgG (γ chain specific)储存液含50%甘油和50%PBS, pH7.4, 无防腐剂。保存于2-8°C; 若需长期保存, 请置于-20°C冰箱。

注意事项

试验中请穿着试验服并带手套做好防护工作。请按实验室安全操作规范进行实验。

本试剂仅供科研使用, 请勿用于临床诊断或其他治疗用途。

部分文献引用

1. Wu Y, Quan Y, Liu Y, et al. Hyperglycaemia inhibits REG3A expression to exacerbate TLR3-mediated skin inflammation in diabetes [J]. Nature Communications, 2016.
2. Wang S, Xu M, Li X, et al. Exosomes released by hepatocarcinoma cells endow adipocytes with tumor-promoting properties [J]. Journal of hematology & oncology, 2018.
3. Liang X, Zhang L, Wang S, et al. Exosomes secreted by mesenchymal stem cells promote endothelial cell angiogenesis by transferring miR-125a [J]. J Cell Sci, 2016.
4. Li H, Fan J, Fan L, et al. miRNA-10b Reciprocally Stimulates Osteogenesis and Inhibits Adipogenesis Partly Through the TGF- β /SMAD2 Signaling Pathway[J]. Aging and Disease.
5. Song G, Chen C, Wu Q, et al. Selenium-enriched yeast inhibited β -amyloid production and modulated autophagy in a triple transgenic mouse model of Alzheimer's Disease [J]. Metallomics, 2018.
6. Li X, Wang S, Zhu R, et al. Lung tumor exosomes induce a pro-inflammatory phenotype in mesenchymal stem cells via NF κ B-TLR signaling pathway [J].Journal of Hematology & Oncology, 2016.
7. Zhang Z H, Wu Q Y, Zheng R, et al. Selenomethionine mitigates cognitive decline by targeting both tau hyperphosphorylation and autophagic clearance in an Alzheimer's disease mouse model[J]. Journal of Neuroscience, 2017.

技术支持

tech@neobioscience.com

全国统一客服

4006-800-892

订货热线

0755-26755892 (深圳)

010-88594029 (北京)

021-34613729 (上海)

18028025014 (广州)